

Food and Drug Administration, HHS

§ 522.1010

and other respiratory problems should exercise extreme caution when handling this product. In the early stages, women may be unaware of their pregnancies. Fluprostenol is readily absorbed through the skin and can cause abortion and/or bronchiospasm. Direct contact with the skin should therefore be avoided. Accidental spillage on the skin should be washed off immediately with soap and water.

[44 FR 52191, Sept. 7, 1979, as amended at 47 FR 22092, May 21, 1982]

§ 522.1002 Follicle stimulating hormone.

(a)(1) *Specifications.* Each package contains 2 vials. One vial contains dry, powdered, porcine pituitary gland equivalent to 75 units (NIH-FSH-S1) of follicle stimulating hormone. The other vial contains 10 milliliters of aqueous diluent.

(2) *Sponsor.* See No. 052923 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) *Dosage.* 12.5 units of follicle stimulating hormone twice a day for 3 days (a total of 75 units). To effect regression of the corpus luteum, prostaglandin should be given with the 5th dose.

(ii) *Indications for use.* For induction of superovulation in cows for procedures requiring the production of multiple ova at a single estrus.

(iii) *Limitations.* For intramuscular use in cows that are not pregnant and have a normal corpus luteum. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(b)(1) *Specifications.* The drug is a lyophilized pituitary extract material. Each 10-milliliter vial contains an amount equivalent to 50 milligrams of standard porcine follicle stimulating hormone and is reconstituted for use by addition of 10 milliliters of 0.9 percent aqueous sodium chloride solution.

(2) *Sponsor.* See 063112 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) *Dosage.* Cattle and horses, 10–50 milligrams; sheep and swine, 5–25 milligrams; dogs, 5–15 milligrams.

(ii) *Indications for use.* The drug is used as a supplemental source of follicle stimulating hormone where there is a general deficiency in cattle, horses, sheep, swine, and dogs.

(iii) *Limitations.* Administer intramuscularly, subcutaneously, or intravenously. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 47377, Sept. 9, 1993, as amended at 62 FR 62242, Nov. 21, 1997; 76 FR 2808, Jan. 18, 2011]

§ 522.1004 Fomepizole.

(a) *Specifications.* Each vial contains 1.5 grams fomepizole (1.5 milliliter (mL) of 1.0 gram per mL solution).

(b) *Sponsors.* See Nos. 046129 and 063286 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount.* 20 milligrams per kilogram (mg/kg) of body weight intravenously initially, followed by 15 mg/kg at 12 and 24 hours, and 5 mg/kg at 36 hours.

(2) *Indications for use.* As an antidote for ethylene glycol (antifreeze) poisoning in dogs who have ingested or are suspected of having ingested ethylene glycol.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[61 FR 68147, Dec. 27, 1996, as amended at 71 FR 28266, May 16, 2006; 74 FR 26952, June 5, 2009; 74 FR 47725, Sept. 17, 2009; 77 FR 26697, May 7, 2012]

§ 522.1010 Furosemide.

(a) *Specifications*—(1) Each milliliter (mL) of solution contains 50 milligrams (mg) furosemide monoethanolamine.

(2) Each mL of solution contains 50 mg furosemide diethanolamine.

(b) *Sponsors.* See sponsors in § 510.600(c) of this chapter for use of products described in paragraph (a) of this section for use as in paragraph (d) of this section.

(1) No. 000010 as described in paragraph (a)(1) of this section for use as in paragraphs (d)(1) and (d)(2)(ii) of this section.

(2) No. 061623 as described in paragraph (a)(2) of this section for use as in paragraph (d)(2)(ii) of this section.

(3) No. 000859 as described in paragraph (a)(2) for use as in paragraphs (d)(1), (d)(2)(i), and (d)(3) of this section.

(4) No. 000061 as described in paragraph (a)(2) for use as in paragraphs (d)(1), (d)(2)(iii), and (d)(3) of this section.